

Parasitic infestations in houseflies and soils collected from garbage dumps in parts of Ijebu-North, Ogun State Nigeria: A cross-sectional study

Adekunle, O.N¹, Mogaji, H.O^{2,3}, Adeleke, M.T¹, Adesetan, T.O⁴, Abimbola-Okuneye W.A, Ogunniran, D.O¹, Ogunade, A.O.¹ and Agbolade, O.M¹

¹Department of Zoology and Environmental Biology, ⁴Department of Microbiology, Olabisi Onabanjo University, Ago-Iwoye Ogun State, Nigeria

²Department of Animal and Environmental Biology, Federal University Oye-Ekiti, Ekiti State, Nigeria

³Instituto of Collective Health, Universidade Federal de Bahia, Brazil

E-mail: adekunle.oladunni@oouaqoiwoye.edu.ng

ABSTRACT

Open defecation and indiscriminate disposal of waste is a common practice and has been highlighted as a risk factor promoting transmission of gastrointestinal parasites hence, we investigated the prevalence of parasites in Houseflies (*Musca domestica*) and soil samples from garbage dumps in Ijebu-North, Ogun State, Nigeria, where such data is non-existent. Flies were collected using baited traps and approximately 100g of surface soil were collected from the garbage dumps in the study area. Standard flotation and sedimentation techniques were used to process collected soil samples, including external and internal contents from *Musca domestica* (*M. domestica*). Isolated parasites were identified under a binocular microscope. Our findings showed that all the houseflies examined were infested with at least one kind of parasite on either external or internal surfaces. A total of 328 (78.8%) of the flies were infested with *Taenia* sp., followed by 296 (74.0%) for *Ascaris lumbricoides*, 200 (48.1%) for hookworms, and the least was 64 (15.6%) for *Toxocara canis*. There were significant differences in the proportions observed across the different garbage dumps in the study area for Hookworm ($p=0.048$), *Fasciola* sp. ($p=0.002$) and *Toxocara canis* ($p=0.033$). On their internal surfaces by location, Oru had the highest prevalence for all the parasites except for Hookworms, *Toxocara canis*, *Strongyloides stercoralis* and *Trichuris trichuria*. Significant differences were observed across the garbage dumps in the study area and Hookworm ($p = 0.015$) and *Strongyloides stercoralis* ($p = 0.014$). Similarly, at least one kind of parasite was recovered from all the 32 soil samples examined, with *Taenia* sp. (90.0%), *Ascaris lumbricoides*. (83.3%) and Hookworms (60.0%) been the most common parasites. This study highlights an important issue of gross environmental contamination with parasite eggs/larvae across garbage dumps which requires prompt clearing of garbage dumps and proper sanitization to prevent disease epidemics.

Keywords: Garbage dumps, *Musca domestica*, Nigeria, Parasites, Soil

Received: 01.03.23

Accepted: 03.07.23

1 INTRODUCTION

Open defecation and indiscriminate disposal of waste is a common practice in marginalized settings, and has been highlighted as a risk factor promoting transmission of gastrointestinal parasites (WHO/UNICEF, 2021). Garbage dumps consisting of dumped food wastes, papers, plastic packaging and metals provide breeding grounds for important vectors of communicable diseases such as houseflies and rodents (Iboh et al., 2015; Ike-Ihunwo

and Gboeloh, 2019; Haywood et al., 2021). Houseflies have been implicated in the transmission cycle of more than 100 pathogens including gastrointestinal parasites which are known to cause a range of morbidities including anemia, malnutrition, lowered cognition, amongst others (Elkanah et al., 2020). Houseflies are known for their ubiquitous presence and preference for breeding in unhygienic surroundings, such as areas contaminated with human or animal

excreta, carcasses, organic waste, and exposed garbage, As such, they continue to serve as a notable mechanical vector for the transmission of pathogens (Nwangwu et al., 2013). The structural adaptations on the external and internal body surfaces of houseflies are suited for picking dirt including ova stages of parasites from garbage dumps and further transmitting them (Gordon and Lavoipierre, 1976; Awache and Farouk, 2016; Farghly et al., 2016). There is plethora of evidence from Nigeria on the prevalence of parasites in houseflies (Adenusi and Adewoga, 2013; Oghale et al., 2013; Balla et al., 2014; Amaechi et al., 2017; Deakpe et al., 2018; Khamesipour et al., 2018; Nwadike and Agbolade, 2020; Elkanah et al., 2020), and on soil samples (Adewole and Ajayi, 2010; Lebari 2021).

However, the degree of parasite contamination in houseflies and soils from garbage dumps is largely unknown in some parts of the country, for instance Ogun State. An available study reported the high prevalence of houseflies from several garbage dumps in Ogun State, with no information on parasites infestation (Banjo et al., 2012). It is therefore important to provide up-to-date data on parasite infestation in houseflies and soil samples to support efforts targeted at controlling

disease epidemic (Oyebamiji et al., 2018). This present study therefore aimed to provide information on parasite infestations in houseflies and soil samples collected from garbage dumps across three communities in Ijebu-North Local Government Area, Ogun State, Nigeria.

2 MATERIALS AND METHODS

2.1 Study Area and Design

This study was conducted between August and December 2021, employing a cross-sectional design and purposive selection of 16 garbage dumps across three communities (Ago-Iwoye, Oru and Ijebu-Igbo) located in Ijebu-North a Local Government Area (LGA) within the study area (Figure 1). Ogun State, Southwest Nigeria at $6^{\circ} 57'N$ $4^{\circ}00'E$ has an area of 967 km^2 with a population of 420,817 (2021 NPC estimated census). The study area is located in the tropical rain forest belt with hot and humid climatic conditions. The average temperature is $27.6 \text{ }^{\circ}\text{C}$ with mean annual precipitation of 2020mm. The indigenes of these towns are mainly Ijebus (Yoruba) with a relatively little proportion of non-indigenous people. It is also a host to a number of higher educational institutions.




Figure 1: Map showing the visited garbage dumps in Ijebu-North Local Government Area. This Figure was created by the authors in ArcGIS 9.3 Software using their primary data and base shapefiles from openly accessible site. The authors therefore give permission for re-use of this map.

2.3 Processing of Samples

The internal and external surfaces of the trapped *M. domestica* were processed using sedimentation techniques as described by Nwadike and Agbolade (2020). The housefly was first introduced into 5mL normal saline (0.85% NaCl solution) in a test tube, shaken less vigorously to disallow disintegration of body parts, and the resultant saline was concentrated by centrifugation at 2000 rpm for 5 minutes. The flies were later dissected under a microscope and the intestinal content was centrifuged at 2000 rpm for 5 minutes. The resulting precipitate from each procedure was stained with 1 drop of Lugol's iodine. Sucrose floatation and Baermann technique was employed in isolation of parasites eggs, larvae or cysts from the soil samples. This method has been previously described by Nkouayep et al. (2017) and Imalele et al. (2021). A trichome staining procedure was used to ascertain *Entamoeba histolytica* (*E. histolytica*). All prepared slides were observed under a binocular microscope using x10 and x40 objective lens.

The Atlas of Medical Helminthology and Protozoology by Sullivan (2009) was used as a reference guide to identify isolated eggs, larvae or cysts.

2.4 Statistical Analysis

Using Social Packages for Social Sciences (version 25) for analysis, descriptive statistics: frequencies and percentages were employed, followed by chi-square tests to estimate the degree of association between parasites prevalence and location. A significant difference was established when $p \leq 0.05$.

3 RESULTS

3.1 Overview of houseflies collected and Examined

(Table 1) were collected across the three communities in Ijebu-North LGA; majority 152 (36.6%) of the houseflies were collected from Ago-Iwoye, followed by 144 (34.6%) from Ijebu-Igbo and 120 (28.8%) from Oru. All the *M. domestica* collected parasites, except in Ago-Iwoye, where only 136 of the 152 houseflies were examined for internal parasites. were examined for external and internal prevalence was recorded for *Toxocara canis* (*T. canis*) eggs 64 (15.4%). By location, Ijebu-Igbo had the highest prevalence for *A. lumbricoides*, *E. histolytica* cysts and *T. canis*. There were significant differences in the prevalence of hookworms, *Fasciola* sp. and *T. canis* across the study communities ($p < 0.05$).

Table 1: Overview of houseflies collected and examined

Communities	Number of houseflies examined for external parasites	Number of houseflies infected with at least one external parasite (%)	Number of houseflies examined for internal parasites	Number of houseflies infected with at least one internal parasite (%)
Ago Iwoye	152	152 (100)	136	136 (100)
Oru	120	120 (100)	120	120 (100)
Ijebu-Igbo	144	144 (100)	144	144 (100)
Total	416	416 (100)	400	400 (100)

*: houseflies infested with at least one kind of parasite; *S. stercoralis* - *Strongyloides stercoralis*; NE: Number Examined; NI: Number Infected

3.2 Prevalence of parasites on the external surface of *M. domestica*

As shown in Table 2 below, all the 416 *M. domestica* examined had at least one kind of parasites on their external surface. An overall

prevalence of 100% was observed. By species', prevalence was highest for eggs of *Taenia* sp. 328 (78.8%), followed by *Ascaris lumbricoides* (*A. lumbricoides*) eggs 296

(71.2%), Hookworms 200 (48.1%), *Fasciola* sp. 192 (47.1%) and *Trichuris trichiuria* (*T. trichiuria*) eggs 192 (46.2%). The lowest

Table 2: Prevalence of parasites on the external surface of *M. domestica* examined

Parasites	COMMUNITIES				p-value
	Ago-Iwoye (NE=152)	Ijebu-Igbo (NE=144)	Oru (NE=120)	Total (N=416)	
	NI (%)	NI (%)	NI (%)	NI (%)	
Infested houseflies*	152 (100)	144 (100)	120 (100)	416 (100)	
<i>Taenia</i> sp.	112(73.7)	112 (77.8)	104 (86.7)	328 (78.8)	0.649
<i>Ascaris lumbricoides</i>	96 (63.2)	112 (77.8)	88 (73.3)	296 (71.2)	0.603
Hookworms	40 (26.3)	80 (55.6)	80 (66.7)	200 (48.1)	0.048
<i>Fasciola</i> sp.	24 (15.8)	80 (55.6)	88 (73.3)	192 (47.1)	0.002
<i>Trichuris trichiuria</i>	72 (47.9)	56 (38.9)	72 (53.3)	192 (46.2)	0.703
<i>Entamoeba histolytica</i>	40 (26.3)	80 (55.6)	24 (20.0)	144 (34.6)	0.065
<i>S. stercoralis</i>	24 (15.8)	48 (33.3)	56 (46.7)	128 (30.8)	0.147
<i>Toxocara canis</i>	8 (5.3)	48 (33.3)	8 (6.7)	64 (15.6)	0.033

*: houseflies infested with at least one kind of parasite; *S. stercoralis* - *Strongyloides stercoralis*; NE: Number Examined; NI: Number Infected

3.3 Prevalence of parasites on the internal surface of *M. domestica* examined

All the houseflies examined were infested with at least one kind of parasite on their internal surface (Table 3). By species', a total of 296 (74.0%) of the flies were infested with *A. lumbricoides* (74.0%), 296 (74.0%) for *Taenia* sp., followed by *Fasciola* sp. with 232 (58.0%), 208 (52.0%) for Hookworms and

192 (48.0%) for *E. histolytica*. The lowest prevalence was recorded with 8 houseflies (2.0%) infested with *Enterobius vermicularis* (*E. vermicularis*) cysts. By location, Ago-Iwoye had the least prevalence for all the parasites except for *Fasciola* sp., *T. canis*, *Strongyloides stercoralis* (*S. stercoralis*) larvae and *T. trichiuria*. There were significant differences in the prevalence of *Taenia* sp., Hookworms and *S. stercoralis* across the study communities ($p < 0.05$).

Table 3: Prevalence of parasites on the internal surface of houseflies examined

Parasites	COMMUNITIES				p-value
	Ago-Iwoye (NE=136)	Ijebu-Igbo (NE=144)	Oru (NE=120)	Total (N=400)	
	NI (%)	NI (%)	NI (%)	NI (%)	
Infested houseflies*	136 (100)	144 (100)	120 (100)	400 (100)	
<i>Ascaris lumbricoides</i>	88 (64.7)	112 (77.8)	96 (80.0)	296 (74.0)	0.555
<i>Taenia</i> sp.	72 (52.9)	120 (83.3)	104 (86.7)	296 (74.0)	0.050
<i>Fasciola</i> sp.	88 (64.7)	56 (38.9)	88 (73.3)	232 (58.0)	0.108
Hookworms	40 (29.4)	112 (77.8)	56 (46.7)	208 (52.0)	0.015
<i>Entamoeba histolytica</i>	56 (41.2)	72 (50.0)	64 (53.3)	192 (48.0)	0.772
<i>Trichuris trichiuria</i>	48 (35.3)	88 (61.1)	40 (33.3)	176 (44.0)	0.187
<i>S. stercoralis</i>	40 (29.4)	88 (61.1)	16 (13.3)	144 (36.0)	0.014
<i>Toxocara canis</i>	32 (23.5)	24 (16.7)	0 (0)	56 (14.0)	0.147
<i>Enterobius vermicularis</i>	0(0)	0(0)	8 (6.7)	8 (2.0)	0.304

NE: Number Examined; NI: Number Infected

*: Soil samples infested with at least one kind of parasite; *S. stercoralis* - *Strongyloides stercoralis*;

NE: Number Examined; NI: Number Infected

3.4 Overview and Prevalence of Parasites on Soil Samples collected and examined

As shown in Table 4, a total of 32 soil samples were examined across the three communities in Ijebu-North LGA; majority of the samples 12 (37.5%) were collected from Ijebu-Igbo, followed by 10 (31.25%) from Oru and 10 (31.25%) from Ago-Iwoye. All the 32 soil samples examined had at least one kind of parasites. An overall prevalence of 100% was observed. Prevalence was highest with 24 (75.0%) soils infested with *Taenia* sp., followed with 22 (68.8%) for *A. lumbricoides*, *Fasciola* sp. 18 (56.2%) and Hookworms 14 (43.8%).

The lowest prevalence in the soils was recorded for *E.vermicularis* and *T. trichuria* with 5 (15.6%) and 5 (15.6%) respectively. By location, *A. lumbricoides* (83.3%), *S. stercoralis* (33.3%), *T. canis* (33.3%) and *E. vermicularis* (25.0%) were more common in soils collected from Ijebu-Igbo (83.3%). *Taenia* sp. (90.0%), *Fasciola* sp. (70.0%) and *T. Trichuria* (20.0%) was more common in Ago-Iwoye. However, Hookworms (60.0%) and *E. histolytica* (40.0%) was more common in Oru. There were no significant differences in the prevalence of parasites recovered from soil samples across the study communities ($p > 0.05$).

Table 4: Prevalence of parasites on soil samples examined

Parasites	Communities				p-value
	Ago-Iwoye (NE=10)	Ijebu-Igbo (NE=12)	Oru (NE=10)	Total (N=32)	
	NI (%)	NI (%)	NI (%)	NI (%)	
<i>Infested soil samples*</i>	10(100)	12 (100)	10(100)	32(100)	
<i>Taenia</i> sp.	9 (90.0)	8(66.7%)	7 (70.0)	24 (75.0)	0.411
<i>Ascaris lumbricoides</i>	4 (40.0)	10 (83.3)	8 (80.0)	22 (68.8)	0.060
<i>Fasciola</i> sp.	7 (70.0)	5 (41.7)	6 (60.0)	18 (56.2)	0.394
Hookworm	3 (30.0)	5 (41.7)	6 (60.0)	14 (43.8)	0.394
<i>Entamoeba histolytica</i>	3 (30.0)	3 (25.0)	4 (40.0)	10 (31.2)	0.748
<i>Toxocara canis</i>	2 (20.0)	4 (33.3)	2 (20.0)	8 (25.0)	0.701
<i>S. stercoralis</i>	1 (10.0)	4 (33.3)	1(10.0)	6 (18.8)	0.262
<i>Enterobius vermicularis</i>	1 (10.0)	3 (25.0)	1 (10.0)	5 (15.6)	0.527
<i>Trichuris trichiuria</i>	2 (20.0)	2 (16.7)	1 (10.0)	5 (15.6)	0.821

*: Soil samples infested with at least one kind of parasite; *S. stercoralis* - *Strongyloides stercoralis*; NE: Number Examined; NI: Number Infected

4 DISCUSSION

Our study provided information on parasites associated with houseflies and soil samples collected from different garbage dumpsites in parts of Ijebu-North, Ogun State, Nigeria. Houseflies are biologically adapted to efficiently transfer pathogenic organisms from their body surface and gut to humans. They accomplish this by depositing these organisms on human food through their activities, making them effective mechanical vectors for disease transmission (Deakpe et al., 2018). Our results shows that all the captured houseflies were infected with at least one parasite either on their external or internal surface. This is consistent with report by Mbakwe et al. (2021) which noted that infectious disease-causing organisms frequently adhere to the mouth, body, hairs and sticky pads of the feet of these flies. The results also indicate that all the houseflies caught in the study area have at least a 50% chance of being exposed to a garbage dump infected with faecal parasites. *M. domestica* is closely associated with faeces, as the latter serves as a protein source and substrate for

the growth of the former (Adenusi and Adewoga, 2013). The low-earning population who cannot afford good housing structures with potable good water and toilets may be forced to make use of their surroundings including garbage dumpsites for the passing of faeces and urine. These practices have been previously highlighted in previous studies (Iboh et al., 2015; Ahmadu et al., 2016; Lebari, 2021).

Our findings are also consistent with other published reports, where parasites recovered from the external surfaces of the examined houseflies were also detected in the internal guts of the flies except for *E. vermicularis* (Deakpe et al., 2018; Issa, 2019; Szalanski et al., 2004). These findings suggests that faeces and vomits may also serve as major routes of transmission of pathogens (Khamesipour et al., 2018). Furthermore, we found very high prevalence of *Taenia* sp. eggs on the external surface (78.8%) and internal surface (74%) of the flies examined. The findings are particularly higher than previous reports by Nwadike and Agbolade (2020), Adenusi and Adewoga (2013) and

Ahmadu et al. (2016). These differences could be attributed to contamination of the houseflies with faecal droppings of high numbers of straying animals (cows and pigs) in the vicinity. The presence of *T. canis* in the results suggests the presence of infested dogs in the study area. Similarly, *A. lumbricoides*, Hookworm, *T. trichiura* and *E. histolytica* which are generally known to cause gastrointestinal diseases in humans, and are significantly associated with morbidities like anaemia, malnutrition, and lower cognitive abilities were recovered. These parasites have also being listed by Khamesipour et al. (2018).

On the other-hand, our findings show significant similarities in the range of parasites recovered across the study sites which reiterated that residents irrespective of their location may likely share similar risk that contributes to the preponderance of these gastrointestinal parasites. Although, this present study did not investigate the potential risk factors associate with parasite infestation, common factors from literatures includes practice of open defecation, unmanaged animal waste near animal or human activities, environmental and seasonality conditions for insect vector proliferation and ingestion of infected insect vectors by animals or humans (Patel et al., 2022). Our analysis of soil sample, corroborates the evidence of soil contamination with infective eggs, which could be transmitted to inhabitant who do not practice adequate hand-mouth hygiene, or do-not engage in regular usage of footwear. Other studies within the country, and elsewhere have extensively reported the contamination of soils from garbage sites with pathogenic protozoan and helminths. (Ikpeama et al., 2016; Adesewa and Morenikeji, 2017; Nkengazong et al., 2021; Steinbaum et al., 2016). Hence it is important to address social-infrastructure disparities and implement interventions targeted at modifying behavioral practices such as open defecation, and poor hand to mouth hygiene that increase exposure to these pathogens.

4.1 CONCLUSION

In conclusion, the transmission of gastrointestinal parasites could be associated with the houseflies and soils gotten from the garbage dumps in the study area. Constant health enlightenment programs should be done on quarterly basis on the implication of open defecation and improper disposal of wastes, especially in relation to disease transmission with emphasis on houseflies as an agent of transmission. Appropriate usage of footwears, hand-washing hygiene with soaps, improved food preparation/processing, and regular deworming exercise should also be prioritized in developed enlightenment programs.

ACKNOWLEDGMENTS

We would like to thank all people in the community who consented to us collecting houseflies and soil from the garbage dumps and also Mr. A. Subair of the Zoology Laboratory of OOU, Ago-Iwoye for his supervision in the Laboratory.

REFERENCES

- Adenusi, A.A. and Adewoga, T.O. 2013. Studies on the potential and public health importance of non-biting synanthropic flies in the mechanical transmission of human enterohelminths. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 107: 812–818.
- Adesewa, A. and Morenikeji, O. 2017. Helminths and heavy metals in soils from dumpsites in Ibadan city, Nigeria. *Journal of Preventive Medicine and Hygiene*, 58: E328-333.
- Adewole, S.O. and Ajayi, A.A. 2010. Studies on human intestinal parasites in refuse dumps. *Journal of Life Sciences*, 4(1): 55-59.
- Ahmadu, Y.M., Goselle, O.N., Ejimadu, L.C. and James-Rugu, N.N. 2016. Microhabitats and pathogens of houseflies (*Musca domestica*): Public health concern. *Electronic Journal of Biology*, 12(4): 374-380.
- Amaechi, A.A., Ukaga, C.N., Iwunze, J.I., Nwachukwu, M.O. and Anumudu, B. 2017. Epidemiological implications of houseflies (*Musca domestica*) in the dessimination of diseases in Owerri, south-east Nigeria. *Nigerian Journal of Parasitology*, 38(2): 298-301.

- Awache, I. and Farouk, A.A. 2016. Bacteria and fungi associated with houseflies collected from cafeteria and food Centres in Sokoto. *FUW Trends in Science and Technology Journal*, 1(1):123-125.
- Balla, H.J., Usman, Y. and Muhammad, A. 2014. The role of housefly (*Musca domestica*) in mechanical transmission of intestinal parasites in Maiduguri metropolis, North Eastern Nigeria. *Journal of Natural Sciences Research*, 4(8):www.iiste.org. ISBN 2224-3186.
- Banjo, F.M., Banjo, A.D. and Fasunwon, B.T. 2012. Survey of arthropod associated with refuse disposal sites in Ijebu-Ode, Ogun State. *Current Research Journal of Biological Sciences*, 4(4): 381-384.
- Dada E. O. and Egbunu A. A. 2016. Dispersion of human intestinal Geohelminth ova in selected refuse dumpsites in Igbara-Oke, Ifedore, Local Government Area, Ondo State, Nigeria. *International Journal of Current Microbiology and Applied Sciences*, 5(4): 924-928.
- Deakpe, T.E., Manyi, M.M. and Utume, L.N. 2018. Pathogenic parasites and bacteria associated with the housefly (*Musca domestica*) in Makurdi; a fly-infested area in central Nigeria. *Nigerian Journal of Parasitology*, 39(1): 111-115.
- Elkanah, S. O., Elkanah, D. S., Fumlack, L. D., Wahedi, J. A. and Kela, S. L. 2020. Houseflies (*Musca Domestica*) as potential carriers of human intestinal parasites in Jalingo Metropolis. *FUDMA Journal of Sciences*, 4(3): 402-405.
- Farghly, A.M., Mohamed, S.M., Abdel-Rahman, S.A., Mohammed, F.E., El-Bahaie, E.S. and El-Shafey, M.A. 2016. The relation between the prevalence of soil-transmitted parasites in the soil and among school children in Zagazig district, Sharkya Governorate, Egypt. *Journal of Parasitic Diseases*, 40(3): 1021-1029.
- Gordon, R.M. and Lavoipierre, M.M.J. 1976. Entomology for students of medicine. (Blackwell Scientific Publication, Oxford, London, Edinburgh & Melbourne, 1976).
- Harroff, L.A., Liotta, J.L., Bowman, D.D. and Angenent, L.T. 2019. Current time-temperature relationships for thermal inactivation of Ascaris eggs at mesophilic temperatures are too conservative and may hamper development of simple, but effective sanitation. *Water Research X*, 5: 10-36.
- Haywood, L.K., Kapwata, T., Oelofse, S., Breetzke, G. and YaelWright, C. 2021. Waste disposal practices in low-income settlements of South Africa. *International Journal of Environmental and Research and Public Health*, 18(15): 8176.
- Iboh, C.I., Effiom, O.E., Ekor, S.C. and Inyang, A. 2015. Parasitological evaluation of un- disposed refuse dumps in Calabar South, Cross River State, Nigeria. *International Journal of Pure and Applied Zoology*, 3(3): 232-239.
- Oyebamiji, D.A., Ebisike, A.N., Egede, J.O. and Hassan, A.A. 2018. Knowledge, attitude and practice with respect to soil contamination by soil-transmitted helminths in Ibadan, Ike-Ihunwo, C. N. and Gboeloh, L.B. 2019. Prevalence of vectors of public health importance in major dumpsites in Port Harcourt Metropolis, Rivers State, Nigeria. *South Asian Journal of Parasitology*, 2(4): 1-9.
- Ikpeama, C., Obianjuru, I. O. C. and Igomaka, A. I. 2016. The impact of refuse disposal dumpsites on the spread of intestinal helminthiasis in Owerri Metropolis, Imo state, South Eastern Nigeria. *International Journal of Clinical & Laboratory Medicine*, 2(2): 13-18.
- Imalele, E.E., Offiong, E.E., Ukam, U.A., Urimaneh, A.W. and Utibe, H.V. 2021. Occurrence of soil-transmitted helminths from selected dumpsites and farmlands in Calabar, Nigeria. *Annual Research & Review in Biology*, 36(5): 14-22.
- Issa, R. 2018. *Musca domestica* acts as transport vector hosts. *Bulletin of the National Research Centre*, 43: 73.
- Lebari, B. G. 2021. Evaluation of gastrointestinal parasites in major dumpsites and health risk behaviour of scavengers in Port Harcourt Metropolis. *EAS Journal of Parasitology and Infectious Diseases*, 3(4): 44-53.
- Mbakwe, O.N., Iwueze, M.O., Irikannu, K.C. and Aniefuna, C.O. 2021. Parasites harbored by synanthropic flies in urban and rural areas of Anambra State, Southeastern Nigeria. *South Asian Journal of Parasitology*, 5(4): 22-28.
- Nkengazong, L., Djabidatou O., Kame, N.G., Irma, A., Efietngab, T.F., Jean, B., Ngue, M. and Zebaze, T.S. 2021. Impact of Hygiene and Sanitation in Cameroon (HYSACAM) refuse dumpsites on the propagation of resistant forms of parasites in the city of Yaoundé, Centre Region-Cameroon. *International Journal of Environmental and Research and Public Health*, 8 (3): 159-207.
- Nkouayep, V.R., Tchakounté, B.N. and Poné, J.W. 2017. Profile of Geohelminth Eggs, Cysts, and Oocysts of Protozoans Contaminating the Soils of Ten Primary Schools in Dschang, West Cameroon. *Journal of Parasitology Research*. Article ID 1534675.
- Nwadike, F.U. and Agbolade, O.M. 2020. Parasites associated with houseflies from some dump sites in Ilisan and Ikenne, Ogun State, Southwest Nigeria. *European Journal of Pharmaceutical and Medical Research*, 7(1): 61-65.
- Nwangwu, U.C., Onyido, A.E., Egbuche, C.M, Iwueze, M.O., and Ezugbo-Nwobi, I.K. 2013. Parasites associated with wild-caught houseflies in Awka metropolis. *IOSR Journal of Pharmacy and Biological Sciences*, 6(1): 12-19.
- Oghale, O.O., Ebube, C.A. and Oluchi, U.O. 2013. Parasitic load on *Musca domestica* (Diptera: Muscidae) from different synanthropic environments in Umuahia metropolis. *Journal of Public Health and Epidemiology*, 5(8): 309-312.
- Southwestern Nigeria. *Parasite Epidemiology and Control*, 3: e00075.

Patel, A., Jenkins, M., Rhoden, K. and Barnes, A.N. A. 2022. Systematic review of zoonotic enteric parasites carried by flies, cockroaches and dung beetles. *Pathogens*, 11: 90.

Steinbaum, L., Njenga, S.M., Kihara, J., Boehm, A.B., Davis, J., Null, C. and Pickering, A.J. 2016. Soil-transmitted helminth eggs are present in soil at multiple locations within households in rural Kenya. *PLoS One*, 11(6): e0157780.

Sullivan, J.T. 2009. A color atlas of parasitology. 8th Edition, University of San Francisco USA. 158pp.

Szalanski, A.L., Owens, C.B., McKay T. and Steelman, C.D. 2004. Detection of *Campylobacter* and *Escherichia coli* O157:H7 from filth flies by polymerase chain reaction. *Medical and Veterinary Entomology*, 18(3): 241–246.

WHO/UNICEF 2021. Joint Monitoring Programme (JMP) for Water Supply, Sanitation and Hygiene.